



Effect of Dietary Supplementation of *Holarrhena Pubescens* Leaf Powder on the Growth Performance, Carcass Characteristics, Microbial Count and Haemato-Biochemical Parameters of Piglets

Olujimi John Alagbe

Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat

Corresponding Author: Olujimi John dralagbe@outlook.com

ARTICLE INFO

Keywords: Antimicrobial, Blood, Carcass, Piglets, Phyto-constituents, Performance

Received : 5 June

Revised : 19 July

Accepted: 20 August

©2024 Alagbe: This is an open-access article distributed under the terms of the

[Creative Commons Atribusi 4.0 Internasional](#).



ABSTRACT

This 60 days experiment was carried out to ascertain the effect of dietary supplementation of *Holarrhena pubescens* leaf powder (HPLP) on the growth performance, carcass characteristics, intestinal microbial population and some haemato-biochemical parameter of piglets. A total of 50 cross bred male pig with an initial body weight 9.90 ± 0.28 kg were randomly divided into five groups (G_A , G_B , G_C , G_D and G_E) with five replicates (1 animal per replicate). Feed (Standard) used in this study was Corn-soya based diet compounded with Nutritional Research Council (2012) standard. The first week was a pre-feeding period where piglets were acclimatized and dewormed against parasites and provided unlimited accesses to clean water throughout the experimental period. A completely randomized design method was adopted and piglets in G_A was fed standard diet without HPLP while those in G_B , G_C , G_D and G_E received same diet supplemented with HPLP at 30 g, 60 g, 90 g and 120 g in that order. Phytochemical screening of HPLP showed that it contained phenols, terpenoids and flavonoids as the most prominent compounds while others include, alkaloids, tannins, saponins and glycosides. Increase in the supplementation significantly improved ($P<0.05$) average daily weight gain, average daily feed intake, dressing percentage and reduced feed conversion ratio and mortality among animals. *Staphylococcus aureus*, *Escherichia coli*, *Proteus spp*, *Pseudomonas spp* and *Staphylococcus spp* count were significantly suppressed by HPLC, conversely, *Lacobacillus spp* increased across the group

INTRODUCTION

Globally, there is clear demand to reduce antibiotic use in animals and in this way reduce the risk of multi-drug resistance. The use of medicinal plants or herbs have been gaining increasing attention in the livestock industry due to their beneficial effect on feed intake and performance (Alagbe *et al.*, 2020). They have also been reported to be non-toxic, environmental friendly and does not have any withdrawal period (Adewale *et al.*, 2021; Ojediran *et al.*, 2024). Herbs contain phyto-constituents or metabolites (terpenoids, tannins, flavonoids, phenolic compounds etc.), that have been investigated as an effective nutritional strategy to improve the overall health of animals (Musa *et al.*, 2020). Among the potential herbal plant with therapeutic property is *Holarrhena pubescens*.

Holarrhena pubescens is a multipurpose deciduous tree that belongs to the Apocynaceae. It is native to South central China and widely distributed in most parts of Philippines, India, Pakistan, Bangladesh, Thailand, Taiwan and tropical Africa including, Zambia, Zimbabwe, Madagascar, Kenya, Tanzania amongst others (Maroyi *et al.*, 2008). The leaves are oblong and elliptic in shape and the tree can grow up to 1300 m in altitude (Guar *et al.*, 2010).

Ethno-medically, *Holarrhena pubescens* is used in traditional medicine in the treatment of piles, stomach ache, fever, body pain, intestinal parasites, tooth ache, diabetes, dysentery, leprosy, skin infection, gonorrhea, rheumatism and arthritis (Jamadagni *et al.*, 2017). Aqueous extract from the stem bark of *Holarrhena pubescens* are also used to treat snake bite, indigestion, respiratory disorder, malaria, stomach cramps and dysentery (Devi *et al.*, 2013). Ethanolic extract from the roots and stem bark of *Holarrhena pubescens* have been proven to inhibit the activities of pathogenic organisms such as, *Plasmodium falciparum*, *Pseudomonas aeruginosa*, *Escherichia coli* and *Candida albicans* (Dubey *et al.*, 2012).

Previous studies have shown that the use of medicinal plants in piglets have shown that it has growth promoting effects (Maenner *et al.*, 2011; Jacela *et al.*, 2010), influence the intestinal microbiota population (Yan and Kim, 2012), modulation of immune system (Li *et al.*, 2012), enhance the activities of enzymes in the gastro intestinal tract (Yan *et al.*, 2010), scavenge free radicals and thus prevents diseases ((Labaque *et al.*, 2013). Most of these result showed that herbs pose positive effect on the general performance of animals. However, research on the impact of *Holarrhena pubescens* leaf powder on piglets are scanty. This research is important because herbs needs to be supplemented at right dose to get the required result and it will also help to address the elevated reports on antimicrobial resistance.

Therefore, the purpose of this research was to determine the effect of dietary supplementation of *Holarrhena pubescens* leaf powder on the growth performance, carcass characteristics, microbial count and haemato-biochemical parameters of piglets.

LITERATURE REVIEW

Phytochemical screening of the leaves, roots and stem bark has shown that it contains, alkaloids, phenolic compounds, flavonoids, terpenoids, steroids and tannins, glycosides in various concentrations (Dua *et al.*, 2013). Reports have shown that *Holarrhena pubescens* possess numerous pharmacological properties including, anti-inflammatory, antioxidant, antidiarrheal, anti-diabetic, analgesic, anti-urolithic, anti-helminthic, antimicrobial and gastro-protective (Ali *et al.*, 2011).

METHODOLOGY

Description of Experimental Location and Ethical Approval

The study was carried out at the Piggery section, Sumitra Research Institute, Gujarat, India located between $16^{\circ} 20^{\prime}$ N East India between the months of September to November, 2024. All procedures were approved by the ethics committee on Animal Use at the Department of Animal Production in the Institute under protocol number ER/H1D/2-2024.

Holarrhena Pubescens Leaf Collection and Preparation

Freshly harvested *Holarrhena pubescens* leaves were collected from Sumitra Research Institute, Gujarat in August 2023. The identity of the leaves was confirmed at the Taxonomy department of the institute where a specimen was deposited with voucher number TDH/2023TG. The leaves were cleaned and air dried for 10 days in an open shade. The dried leaves were grinded using an electric blending machine and the powdered samples were stored in an air tight plastic container for further laboratory evaluation.

Care of Experimental Animals and Design

A total of 50 cross bred male pigs with an initial body weight of 9.90 ± 0.28 kg and weaned on 28 days were sourced from a reputable breeding farm in Gujarat. Animals were individually housed in pens measuring $1m \times 2m \times 1m$. Prior to the arrival of pigs, pens were thoroughly washed with disinfectants and also disinfected with Morigad Plus®, feeding and watering troughs were properly washed and subsequently cleaned daily. The first week was considered as a pre-feeding stage and pigs were treated against parasites using Oramectin Plus ® dose was administered according to the package insert before they were randomly divided into five groups (A, B, C, D and E) with five replicates (1 animal per replicate). Feed used in this study was Corn-soya based diet compounded with Nutritional Research Council (2012) standard. Feed was provided thrice a day (6:30, 12:00 and 16:30 H). Pigs also had unlimited access to fresh clean water throughout the 60 days trial. A completely randomized design was adopted, animals in group A (control) received standard diet (corn-soya diet) without *Holarrhena pubescens* leaf powder while those in group B, C, D and

E were fed same diet supplemented *Holarrhena pubescens* leaf powder with at 30 g, 60 g, 90 g and 120 g per kilogram diet.

Data Collection

Growth performance parameters such as; average weight gain, average daily weight gain, total feed consumption, average daily feed consumption, feed conversion ratio and percentage mortality was estimated according to methods recently published by John (2024).

Calculations

Average weight gain (kg) = average final body weight - average initial body weight

Average daily weight gain (kg) = average weight gain ÷ 60 (number of experimental period)

Total feed consumption (kg) = feed served - feed refused

Average daily feed consumed (kg) = total feed consumed ÷ 60 (number of experimental period)

Feed conversion ratio = average daily feed consumed ÷ average daily weight gain

Qualitative Phytochemical Analysis of *Holarrhena Pubescens* Leaf Powder

Screening for Alkaloids

One gram of *Holarrhena pubescens* leaf powder was mixed with 10 mL of 1 percent dilute hydrochloric acid on water bath and filtered while hot. Two mL of the filtrate was treated with a few drops of Dragendorff's reagent. An orange brown precipitate was taken as evidence for the presence of alkaloids in the sample (Evans, 2002).

Test for Flavonoids

Five millilitres of dilute ammonia solution was added to one gram of *Holarrhena pubescens* leaf powder followed by addition of concentrated Tetraoxosulphate (VI) acid. A yellow colouration observed indicated the presence of flavonoids. The yellow colouration disappeared on standing (Edeoga et al., 2005).

Test for Tannins

One gram of *Holarrhena pubescens* leaf powder was added to few drops of 1% lead acetate. A yellowish precipitate suggested the presence of tannins (Savithramma et al., 2011).

Test for Saponins

One gram of *Holarrhena pubescens* leaf powder was mixed with 20 mL of distilled water and then agitated for 15 minutes. Formation of foam shows the presence of saponins (Shin et al., 2001).

Test for Glycosides

One gram of *Holarrhena pubescens* leaf powder were treated with 2 mL of glacial acetic acid containing two drops of ferric chloride solution followed by the addition of concentrated sulphuric acid. A reddish brown colouration at the

junction of the two layers and bluish green colour in the upper layer indicated the presence of glycosides (Shin et al., 2001).

Test for Anthraquinones

One gram of *Holarrhena pubescens* leaf powder was added to 5 mL benzene in a test tube and stirred, mixture is allowed to stay for 2 minutes before 10 percent ammonia solution was added. The mixture was shaken and the presence of a red colour suggested the presence of anthraquinones (Evans, 2002).

Test for Anthocyanins

Two grams of *Holarrhena pubescens* leaf powder was added to 5 mL of hydrochloric acid and 2 mL of ammonia. The appearance of pink-red to blue-violet suggest the presence of anthocyanins (Savithramma et al., 2011).

Test for Terpenoids

To 1 g of *Holarrhena pubescens* leaf powder, 1 ml of concentrated sulfuric acid was added and heated to 70 °C for 2 min. A visual check for the development of a grey colouration indicated the presence of terpenoids (Edeoga et al., 2005).

Test for Phenols

To 1 g of *Holarrhena pubescens* leaf powder, 1 ml of 1 % of sodium hydroxide was added and mixed for 1 min. No incubation condition as the colour development is instantaneous. A visual check observing the development of a red colouration indicated the presence of phenols (Edeoga et al., 2005).

Carcass and Organ Weight Evaluation

At the end of the experiment (day 60), five pigs were randomly selected per treatment and feed starved over the night and given access to only clean water. Weights of animals were taken before taking animals to the slaughter slab. Slaughtering procedures followed the method recently published by John (2024). Pigs were first stunned to make them unconscious followed by bleeding, scalding, removal of hair and eviscerated before dissecting into separate parts for carcass and organ measurements. Measurement was carried out on the cut parts which were the head, trotters, back, belly, hams, loins and shoulder. The internal organs such as heart, liver, spleen, kidney, lungs and the large/small intestine were measured with the aid of a digital sensitive scale.

Intestinal Gut Microbial Analysis

At the end of the trial, five pigs were randomly selected per treatment (from dose used for carcass evaluation). One gram of intestinal content was collected into a sterile labeled sample bottles and mixed with 0.1 % peptone saline solution. Collected samples were placed in an ice pack before they were taken immediately to the microbiology laboratory of Sumitra Research Institute, Gujarat, India. Enumeration of each bacterial count was carried out according to recent procedure by Williams et al. (2022) and expressed in Log₁₀ (Cfu).

Blood Collection and Evaluation at the End of the Trial

On the 60th day of the experiment, 5 mL of blood samples were collected from the Jugular vein of five randomly selected rabbits per treatment for blood analysis. 2.5 mL of blood was collected into bottles with anticoagulant for haematological studies while those for serum evaluation was transferred into bottles without anticoagulant before all samples were placed in a prepared ice pack to protect samples before getting to the laboratory. Haematological evaluation was carried out according to methods outlined by Daniel et al. (2024). Commercial diagnostic kit (Kermat automated haematology analyzer; model-LKI -400K, Netherlands). Red blood cell, haemoglobin, pack cell volume and white blood cell counts was determined using flow cytometry differentiation technique (Alagbe, 2023). Same kit was also used to ascertain total albumin, globulin, creatinine, urea, aspartate aminotransferase, alanine aminotransferase and alkaline phosphatase were determined using triangle laser scattering and cyanide free technique. Plasma lipid composition (cholesterol, triglycerides, low density lipoprotein and high density lipoprotein) were accessed using Mac automatic biochemistry analyzer (Model XVB/208D, Netherlands). The machine was calibrated using buffer solutions, thereafter it was adjusted to a sample volume (3.5 - 50.0 μ L), reagent volume (35.0 – 400.0 μ L), wavelength of about 1200 nm and absorbance (0 to 5 Abs).

Laboratory Analysis of Experimental Diet

Analysis of experimental diet was done using commercial kit, NIRTM DA1800 near infra-red feed analyzer which possess the following technical specifications; wavelength range (1200 to 1800 nm), ambient temperature (4 - 45 °C), ambient humidity (< 90 percent relative humidity), and results was generated at an analysis time of less than 1 minutes. Calcium and phosphorus was analyzed using Trace Series Atomic Absorption Spectrometer (Model 12GH/08, Netherlands) following all instructions of the manufacturer to ensure accuracy.

Data Analysis

All the data on growth performance, carcass characteristics, microbial population and blood parameters were subjected to one-way Analysis of Variance using Statistical Package for Social Sciences (version 21). The differences among the treatment means were determined ($P<0.05$) by Duncan multiple range test of the same statistical package.

Table 1. Gross Composition of Experimental Diet Fed to Piglets for 60 Days

Feed items or Ingredients	Content (%)
Corn	53.00
Rice bran	4.00
Soya meal	33.5
Fish meal	2.00
Stone powder (Calcium carbonate)	2.50
Bone meal	4.00
Lysine	0.20
Methionine	0.20
*Premix	0.25
Toxin quarsh binder®	0.01
Common salt	0.34
Total	100.0
Determined analysis	
Crude protein (%)	22.81
Crude fibre (%)	3.61
Crude fat (%)	4.04
Calcium (%)	1.95
Phosphorus (%)	0.68
Metabolizable energy (Kcal/kg)	2700.8

Phytochemical screening of *Holarrhena pubescens* leaf powder (Table 2) revealed that phenols, terpenoids and flavonoids are the most prominent phyto-constituents followed by alkaloids, glycosides, tannins and saponins. Anthracyanins and anthraquinones were absent in the analysis.

Table 2. Phytochemical Screening of *Holarrhena Pubescens* Leaf Powder

Components	Results
Alkaloids	+++
Flavonoids	++++
Terpenoids	++++
Saponins	+
Phenols	++++
Anthracyanins	-
Anthraquinones	-
Tannins	+
Glycosides	+++

+ Present, +++moderately present, +++, highly present - absent

The effect of *Holarrhena pubescens* leaf powder (HPLP) on the growth performance of weaned pigs (Table 3). Average daily body weight was higher ($P<0.05$) for pigs in GE (120 g HPLP/kg diet), intermediate in GB (30 g HPLP/kg diet), GC (60 g HPLP per kg diet) and GD (90 g HPLP per kg diet) as compared to the lowest recorded for GA (control). Average daily feed intake in GB and GC were similar ($P>0.05$) to those pigs which received GD and GE but significantly higher ($P<0.05$) than GA (control). Mortality was higher ($P<0.05$) for GA than the rest of the treatments, with no differences ($P>0.05$) between the other treatments.

Table 3. The Effect of *Holarrhena Pubescens* Leaf Powder on the Growth Performance of Weaned Pigs

Variables	G _A	G _B	G _C	G _D	G _E	SEM	p-value
Number of pigs	15.00	15.00	15.00	15.00	15.00	-	-
Duration of experiment in days	60.00	60.00	60.00	60.00	60.00	-	-
Initial body weight (kg)	10.02	10.10	10.20	10.00	9.99	0.10	0.27
Final body weight (kg)	29.82 ^c	34.03 ^b	34.05 ^b	34.11 ^b	38.92 ^a	0.57	0.34
Body weight gain (kg)	19.80 ^c	23.93 ^b	23.85 ^b	24.11 ^b	28.93 ^a	0.41	0.30
Daily body weight gain (kg)	0.33 ^c	0.40 ^b	0.40 ^b	0.40 ^b	0.48 ^a	0.001	0.0029
Total feed intake (kg)	72.51 ^b	81.23 ^a	82.03 ^a	82.11 ^a	83.04 ^a	1.56	0.61
Daily feed intake (kg)	1.21 ^b	1.36 ^a	1.37 ^a	1.37 ^a	1.38 ^a	0.01	0.28
Feed conversion ratio	3.66 ^a	3.40 ^b	3.41 ^b	3.40 ^b	3.00 ^c	0.01	0.14
Mortality (%)	1.20	-	-	-	-	0.001	0.002

Different letters in the same column (^{a, b, c}) suggests significant differences at $P<0.05$; SEM, standard error of means; each value represents the mean of five replicates (10 pigs per replicate); G_A: basal diet without *Holarrhena pubescens* leaf powder (control; HPLP); G_B: basal diet with 30 g HPLP per kg diet; G_C: basal diet with 60 g HPLP per kg diet; G_D: basal diet with 90 g HPLP per kg diet; G_E: basal diet with 120 g HPLP per kg diet

The effect of *Holarrhena pubescens* leaf powder on the carcass characteristics of weaned pigs (Table 4). Carcass weight was higher ($P<0.05$) in GE (120 g HPLP per kg diet) than GB (30 g HPLP/kg diet), GC (60 g HPLP per kg diet) and GD (90 g HPLP per kg diet) which had similar ($P>0.05$) values while the lowest value was recorded in G_A (control; no HPLP). Dressing percentage values which varied from 62.98 to 70.93 % was higher in GE than other groups. Weights of fore limb, hind limb, belly, back and lungs were higher ($P<0.05$) for GE than GA, but was same ($P>0.05$) with that for the rest of the treatments (G_B, G_C and G_D). Conversely, weights of head, kidney, heart, gastro intestinal tract and liver were not influenced ($P>0.05$) by the treatment.

Table 4. The Effect of *Holarrhena Pubescens* Leaf Powder on the Carcass Characteristics of Weaned Pigs

Variables	G _A	G _B	G _C	G _D	G _E	SEM	p-value
Live weight	29.45 ^c	34.98 ^b	37.12 ^b	37.55 ^b	41.02 ^a	0.67	0.39
Carcass weight	18.55 ^c	23.56 ^b	25.11 ^b	25.04 ^b	29.09 ^a	0.54	0.22
Dressing percentage	62.98 ^c	67.35 ^b	67.65 ^b	66.68 ^b	70.93 ^a	1.75	0.08
Head	2.9	2.91	2.93	2.95	2.99	0.21	0.01
Hind limb	5.88 ^c	7.56 ^b	7.95 ^b	7.98 ^b	8.22 ^a	0.28	0.77
Fore limb	6.94 ^c	8.03 ^b	8.11 ^b	8.13 ^b	8.40 ^a	0.12	0.52
Belly	1.88 ^c	2.14 ^b	2.35 ^b	2.71 ^b	3.00 ^a	0.05	1.26
Back	2.91 ^c	3.09 ^b	3.11 ^b	3.18 ^b	4.01 ^a	0.09	1.40
Liver	0.81	0.85	0.92	0.94	0.97	0.02	0.95
Lungs	0.25 ^c	0.26 ^b	0.27 ^b	0.30 ^b	0.41 ^a	0.01	1.00
Kidney	0.04	0.06	0.07	0.09	0.09	0.02	1.01
Spleen	0.10	0.11	0.11	0.11	0.12	0.01	0.09
Heart	0.15	0.17	0.18	0.20	0.20	0.02	0.08
GIT weight	3.10	3.13	3.17	3.3	3.41	0.74	0.05

Different letters in the same column (^{a, b, c}) suggests significant differences at P<0.05; SEM, standard error of means; each value represents the mean of five replicates (10 pigs per replicate); G_A: basal diet without *Holarrhena pubescens* leaf powder (control; HPLP); G_B: basal diet with 30 g HPLP per kg diet; G_C: basal diet with 60 g HPLP per kg diet; G_D: basal diet with 90 g HPLP per kg diet; G_E: basal diet with 120 g HPLP per kg diet

The effect of *Holarrhena pubescens* leaf powder on intestinal microbial population of weaned pigs (Table 5). Microbial count of *Staphylococcus aureus*, *Escherichia coli*, *Proteus spp*, *Pseudomonas spp* and *Streptococcus spp* was higher (P<0.05) for G_A but was same (P>0.05) with that for the rest of the treatments. Conversely, *Lactobacillus spp* count was similar (P>0.05) among pigs which received G_C, G_D and G_E, intermediate in G_B and lowest in G_A.

Table 5. The Ffect of *Holarrhena Pubescens* Leaf Powder on Intestinal Microbial Population of Weaned Pigs

Microbial population (log10 Cfu)	G _A	G _B	G _C	G _D	G _E	SEM	p-value
<i>Staphylococcus aureus</i>	4.86 ^a	2.92 ^b	2.81 ^b	2.50 ^b	2.31 ^b	0.87	1.12
<i>Escherichia coli</i>	5.77 ^a	3.35 ^b	3.27 ^b	3.16 ^b	3.10 ^b	0.95	1.35
<i>Proteus spp</i>	2.65 ^a	1.95 ^b	1.80 ^b	1.52 ^b	1.41 ^b	0.02	0.46
<i>Pseudomonas spp</i>	2.00 ^a	1.03 ^b	1.00 ^b	0.86 ^c	0.81 ^c	0.01	1.10
<i>Streptococcus spp</i>	3.15 ^a	2.06 ^b	1.95 ^c	1.81 ^c	1.60 ^c	0.03	0.80
<i>Lactobacillus spp</i>	2.07 ^c	3.45 ^b	4.98 ^a	5.02 ^a	5.11 ^a	0.07	1.03

Different letters in the same column (^{a, b, c}) suggests significant differences at P<0.05; SEM, standard error of means; each value represents the mean of five replicates (10 pigs per replicate); G_A: basal diet without *Holarrhena pubescens* leaf powder (control; HPLP); G_B: basal diet with 30 g HPLP per kg diet; G_C: basal diet with 60 g HPLP per kg diet; G_D: basal diet with 90 g HPLP per kg diet; G_E: basal diet with 120 g HPLP per kg diet.

The effect of *Holarrhena pubescens* leaf powder on some haematological indices of weaned pigs (Table 6). Dietary supplementation of *Holarrhena pubescens* leaf powder significantly ($P<0.05$) influenced the hemoglobin, packed cell volume, platelets, red blood cell, white blood cell, lymphocytes, and neutrophil count. On the other hand, there were no significant differences ($P>0.05$) in monocyte count. The pack cell volume percentage, red blood cell, haemoglobin, white blood cell, neutrophils and lymphocytes count was greater ($P<0.05$) for GB, GC, GD and GE as compared to the control (GA).

Table 6. The Effect of *Holarrhena Pubescens* Leaf Powder on Some Haematological Indices of Weaned Pigs

Variables	G _A	G _B	G _C	G _D	G _E	SEM	p-value
Pack cell volume (%)	29.08 ^b	33.56 ^a	34.01 ^a	35.11 ^a	37.09 ^a	1.72	0.09
Haemoglobin (g/L)	87.56 ^c	99.56 ^b	102.7 ^a	109.8 ^a	110.3 ^a	10.94	1.74
Platelet ($\times 10^9/L$)	110.6 ^c	134.7 ^b	139.5 ^a	140.2 ^a	143.9 ^a	11.56	1.11
Red blood cell ($\times 10^{12}/L$)	7.94 ^b	10.65 ^a	10.94 ^a	11.4 ^a	11.6 ^a	0.08	0.03
White blood cell ($\times 10^9/L$)	11.57 ^b	14.87 ^a	15.71 ^a	15.87 ^a	15.76 ^a	0.09	0.02
Neutrophils ($\times 10^9/L$)	6.67 ^{ab}	8.00 ^a	8.05 ^a	9.04 ^a	9.12 ^a	0.43	0.10
Monocytes ($\times 10^9/L$)	0.86	0.81	0.83	0.92	0.95	0.01	0.14
Lymphocytes ($\times 10^9/L$)	9.11 ^b	11.65 ^{ab}	13.86 ^a	14.89 ^a	14.91 ^a	1.20	0.01

Different letters in the same column (^{a, b, c}) suggests significant differences at $P<0.05$; SEM, standard error of means; each value represents the mean of five replicates (10 pigs per replicate); G_A: basal diet without *Holarrhena pubescens* leaf powder (control; HPLP); G_B: basal diet with 30 g HPLP per kg diet; G_C: basal diet with 60 g HPLP per kg diet; G_D: basal diet with 90 g HPLP per kg diet; G_E: basal diet with 120 g HPLP per kg diet

The effect of *Holarrhena pubescens* leaf powder on the serum biochemical indices of weaned pigs (Table 7). Pigs fed diet supplemented with 60 g HPLP per kg diet (G_C), 90 g HPLP per kg diet (G_D), and 120 g HPLP per kg diet (G_E) exhibited higher ($P<0.05$) total protein, albumin, globulin and glucose levels than the other groups. Albumin/globulin ratio (A/G ratio) were similar ($P>0.05$) for pigs which received G_B, G_C, G_D and G_E but significantly higher than those fed G_A (control). On the other hand, activities of alanine aminotransferase, aspartate aminotransferase and alkaline phosphatase were not influenced ($P>0.05$) by the treatments.

Table 7. The Effect of *Holarrhena Pubescens* Leaf Powder on the Serum Biochemical Indices of Weaned Pigs

Variables	G _A	G _B	G _C	G _D	G _E	SEM	p-value
Total protein (g/dL)	5.12 ^c	6.99 ^b	7.09 ^a	7.26 ^a	7.41 ^a	0.04	1.12
Albumin (g/dL)	3.09 ^b	4.12 ^a	4.18 ^a	4.30 ^a	4.42 ^a	0.01	0.03
Globulin (g/dL)	2.03 ^b	2.87 ^a	2.91 ^a	2.96 ^a	2.99 ^a	0.02	0.01
A/G ratio	1.52 ^a	1.43 ^b	1.43 ^b	1.45 ^b	1.47 ^b	0.09	0.07
Glucose (mmol/L)	4.06 ^b	4.56 ^b	6.15 ^a	6.49 ^a	6.59 ^a	2.88	0.05
Alanine aminotransferase (IU/L)	76.87	78.94	79.12	79.15	79.88	9.52	1.00
Aspartate aminotransferase (U/L)	52.75	55.06	55.11	56.03	56.97	4.48	1.06
Alkaline phosphatase (U/L)	102.5	109.6	113.8	115.7	115.9	25.6	7.41

Different letters in the same column (a, b, c) suggests significant differences at P<0.05; SEM, standard error of means; each value represents the mean of five replicates (10 pigs per replicate); G_A: basal diet without *Holarrhena pubescens* leaf powder (control; HPLP); G_B: basal diet with 30 g HPLP per kg diet; G_C: basal diet with 60 g HPLP per kg diet; G_D: basal diet with 90 g HPLP per kg diet; G_E: basal diet with 120 g HPLP per kg diet

The effect of *Holarrhena pubescens* leaf powder on the lipid composition of weaned pigs (Table 8). Cholesterol, triglycerides, low density lipoprotein and high density lipoprotein values were influenced (P<0.05) by the treatments. Pigs fed G_B (30 g HPLP/kg diet), G_C (60 g HPLP per kg diet), G_D (90 g HPLP per kg diet) and G_E (120 g HPLP per kg diet) had a lower cholesterol, triglycerides and low density lipoproteins than the birds fed the control diet (G_A; 0 g HPLP). Conversely, Pigs which received HPLP (G_A, G_B, G_C, G_D and G_E) had a higher values of high density lipoprotein relative to G_A (control).

Table 8: The Effect of *Holarrhena Pubescens* Leaf Powder on The Lipid Composition of Weaned Pigs

Variables (mg/dL)	G _A	G _B	G _C	G _D	G _E	SEM	p-value
Cholesterol	131.6 ^a	110.7 ^b	103.8 ^b	101.7 ^b	100.1 ^b	9.61	1.75
Triglycerides	77.81 ^a	62.17 ^b	60.98 ^b	55.12 ^c	52.09 ^c	3.82	1.20
High density lipoprotein	45.88 ^c	55.71 ^b	58.32 ^b	65.07 ^a	67.05 ^a	4.04	0.53
Low density lipoprotein	110.4 ^a	98.44 ^b	90.31 ^b	87.92 ^c	85.03 ^c	4.75	0.97

Different letters in the same column (a, b, c) suggests significant differences at P<0.05; SEM, standard error of means; each value represents the mean of five replicates (10 pigs per replicate); G_A: basal diet without *Holarrhena pubescens* leaf powder (control; HPLP); G_B: basal diet with 30 g HPLP per kg diet; G_C: basal diet with 60 g HPLP per kg diet; G_D: basal diet with 90 g HPLP per kg diet; G_E: basal diet with 120 g HPLP per kg diet

DISCUSSION

Phytochemical analysis of *Holarrhena pubescens* leaf powder proves the presence of phyto-components or metabolites such as, flavonoids, tannins, alkaloids, phenols, glycosides and terpenoids. Result obtained in this study is in agreement with the study carried out by (Kalimuthu, and Arunprasath, 2018; Kabir et al., 2018). Several of these phyto-components have therapeutic properties and their concentration in the plant tissues is considered as the main factor to evaluate the medicinal value and quality of a given herb (Wills, 2000; Alagbe, 2024). Phenolic compounds in *Holarrhena pubescens* possess a wide array of pharmacological properties, such as antioxidant (Ojediran et al., 2024a), antiulcer, cytoprotective, chemo-preventive, anti-inflammatory, anti-diabetic, antimicrobial, immune-stimulatory, hepatoprotective and free-radical scavenging activity (Ojediran et al., 2024b; Daniel et al., 2023). These attributes are desirable for use of this plant as an agent against the effects that could arise from mycotoxin exposure in humans and animals (Singh et al., 2022). Flavonoids and terpenoids have been suggested to possess gastro-protective and antimicrobial properties, thus it possess excellent inhibitory effects against *Pseudomonas aeruginosa*, *Candida albicans*, *Brevibacillus agri*, *Propionibacterium acnes*, *Trichophyton mentagrophytes*, *Microsporum canis* and *Staphylococcus spp* (Adewale et al., 2021). According to Shittu et al. (2021), tannins performs anti-inflammatory activities and analgesic roles making it a useful therapy in the treatment of yellow fever, malaria, diarrhea, headache, stomach and tooth ache. Medicinally, steroids have been shown to exhibit hormonal, antidiarrhea and anti-Inflammatory activities. They have also been suggested exhibit antifungal, antibacterial, antiviral and hypolipidemic activities (Musa et al., 2020). Saponins can be used as adjuvants during vaccine production and possess a strong gastro-protective property (Muritala et al., 2022). Glycosides have been reported to exhibit wide range of pharmacological activities including, anti-inflammatory and anti-helminthic (Wang et al., 2006; Alagbe, 2021). A synergy in all the phyto-constituents recorded in this experiment could enhance immune response, have strong antibacterial properties, stimulate pancreatic enzyme production and overcome loss of appetite and improve pig's growth performance (Ugbogu and Akukwe, 2009; Alagbe et al., 2021).

Piglets receiving HPLP had an improved average daily weight gain compared to the control, though those fed diet supplemented with HPLP at 120 g per kg diet (G_E) was significantly higher than those which received G_B (30 g HPLP per kg diet), G_C (60 g HPLP per kg diet) and G_D (90 g HPLP per kg diet). This result suggests that higher levels of HPLP may improve nutrient digestion and support the growth of beneficial bacteria as it has an antimicrobial effect and may help protect gut function against the colonization of pathogenic bacteria (John, 2024a; John, 2024b). The average daily weight gain recorded in this study which varied from 0.33 - 0.48 kg was also similar to the result of a study by

(Oetting et al., 2006) who found that average daily weight gain of piglets fed diet supplemented with herbal extracts ranged from 0.21 - 0.55 kg. This result was lower than those presented by (Pastorelli et al., 2006) who recorded an average daily weight gain of 0.21 - 0.32 kg. This variation could be attributed to levels of dietary supplementation, processing method of plants and phyto-constituents in the test material (John, 2024c). Overall, piglets receiving the supplemented feed had a similar feed intake to piglets on the control diet this outcome suggests that HPLP has flavoring effect due to the presence of phyto-constituents, which could arouse an animal's appetite. According to (Best, 2000), Tannins are reported to increase feed palatability, guide the mucous membrane and protect against the risk of having diarrhea or watery fecal droppings in pigs. Flavonoids and phenolic compounds have also shown to demonstrate both antimicrobial and antioxidant functions in the animal and play a role in appetite enhancement (John, 2024d; Tepe et al., 2005). The result obtained on feed intake is in line with the reports of [30] when *Lippia citriodora* was supplemented in the diet of piglets. Feed conversion in the control was 3.66 and decreased to 3.00 in the G_E (120 g HPLP per kg diet). This result indicates that HPLP can promote the transportation of metabolic products and improve feed conversion ratio in pigs resulting in rapid growth promoting (Bartos et al., 2006; Hanezakowska et al., 2012). The result obtained in this study is in agreement with the report of (Hanezakowska et al., 2012) who recorded a feed conversion ratio range of 2.91 - 3.70 when herbal extracts were fed to piglets. Mortality was recorded only among piglets given G_A (control) while none was recorded in the other groups. This result shows that HPLP exhibit antimicrobial, antioxidant, anti-inflammatory properties amongst others (John, 2024e; Taivini et al., 2001). Outcome of this study is in line with the report of (Bartos et al., 2006) when phytogenics were fed to piglets.

Piglets fed a diet containing HPLP increased ($p<0.05$) dressing percentage, fore limb, hind limb, belly, back and lungs compared to those that received the control diet (G_A). Values of dressing percentage (62.98 to 70.93 %), fore limbs (6.94 to 8.13 kg) and hind limbs (5.88 to 8.22 kg) range observed in this experiment was similar to the outcome of a study by [34] who discovered that values obtained varied from 61.00 - 70.04 %, 5.03 - 8.77 kg and 5.11 - 9.00 kg respectively. Weight of lungs was higher in piglets fed G_E compared to the other group, which is likely due to the activities of phyto-constituents in HPLP. The significant increase allows more supply of oxygen needed to drive nutrient round the body of pigs. Weights of head, liver, kidney, spleen, heart and gastro intestinal tract were not influenced ($P>0.05$) by the treatment. This result suggests that the phyto-components in HPLP were not toxic to the pigs (Bartos et al., 2006). Findings of

the present experiment were similar to those recorded by (Czech et al., 2009; Bruno et al., 2013).

Intestinal microbial population of *Staphylococcus aureus* which varied from 2.31 to 4.86 (Log₁₀ Cfu), *Escherichia coli* [(log₁₀ 3.10 to 5.77 Cfu)], *Proteus spp* [(log₁₀ 1.41 to 5.77 Cfu)], *Pseudomonas spp* [(log₁₀ 0.81 to 2.00 Cfu)] and *Streptococcus spp* [(log₁₀ 1.60 to 3.15 Cfu)] decreased in groups fed HLPL compared to that in the control group (G_A). *Lactobacillus spp* population values which ranged from [(log₁₀ 2.07 to 5.11 Cfu)] elevated as the supplementation of HLPL increased in the diet. The outcome of this result indicates that the mode of action of HLPL is based on the principle of competitive exclusion, including eradicating harmful bacteria's (*Staphylococcus aureus*, *Escherichia coli*, *Proteus spp*, *Pseudomonas spp* and *Streptococcus spp*) by producing strong antimicrobial constituents, thus increasing the activities of beneficial bacteria (*Lactobacillus spp*) to mitigate gut health challenges (Bartos et al., 2006). Dietary supplementation of HLPL could also ensure easy absorption of nutrients at an optimum rate and provide efficient protection against pathogens through its own immune system (De Lange et al., 2010).

The present study showed that pack cell volume, platelets, red blood cell and haemoglobin values which varied from 29.08 to 37.09 %, [(110.6 to 143.9 ($\times 10^9/L$)], [(7.96 to 11.6 ($\times 10^{12}/L$))] and [(87.56 to 110.3 g/dL)] was influenced by HPLP supplementation in the diet of piglets. The values obtained in this study is within the baseline values; pack cell volume (20.88 to 40.09 %), platelets [(102.3 to 159.80 ($\times 10^9/L$)], red blood cell ([6.60 - 15.00 ($\times 10^{12}/L$)] and haemoglobin ([95.00 to 152.44 g/dL]) cited by (Winnicka, 2011; Mayengbam and Tolenkhomba, 2015). Haemoglobin is an iron containing protein found in red cells that allows it to carry oxygen (Egeli et al., 1998; Alagbe, 2022). Decrease in pack cell volume and haemoglobin concentration could result in anemia (Faustini et al., 2000; Friendship and Henry, 1996). Though pack cell volume, haemoglobin and red blood cells were higher in groups which received HPLP relative to the control, this result gives room for sufficient oxygen to drive absorbed nutrients round the body of pigs (Chmielowiec et al., 2008; Cooper et al., 2014). Chronic inflammation is one of the causes of low platelet count in pigs, this could affect the overall health of animals (Czech and Grela, 2002; Czech et al., 2009). White blood cell is saddled with the responsibility of producing antibodies to fight against infections and disease in the body (Czech et al., 2009). The lowest white blood cell count was recorded among pigs fed control diet relative to the other groups which is likely to be associated with the increase HPLP supplementation in the diet. The result also suggest that HPLP possess immune-stimulatory and hepatoprotective properties (Shittu et al., 2024). Result in this current study is in agreement with the findings of Chmielowiec et al. (2008). White blood cell

recorded [(11.57 to 15.76 ($\times 10^9/L$))] is within the baseline values for piglets cited by Egeli et al. (1998). Lymphocytes, neutrophils and monocytes values varied from [(9.11 - 14.91 ($\times 10^9/L$))], [(6.67 - 9.12)] and [(0.86 - 0.92 ($\times 10^9/L$))] were within the standard values [(8.77 - 15.80)], 6.00 - 15.00($\times 10^9/L$) and 0.50 - 1.00 ($\times 10^9/L$) reported by Radostits et al. (2000). Lymphocytes and neutrophil values were higher in groups fed diet supplemented with HPLP compared to the control. Lymphocytes and neutrophils are associated with viral and bacterial infections (Radostits et al., 2000). Monocytes play a vital role in maintaining the immune system of pigs (Alagbe, 2024). Macrophages and neutrophils are the leucocytes most used during an inflammation. These blood cells moves to pathogens' entry sites and carry out phagocytosis, eliminating infectious agents. Therefore, HPLP modify the metabolism of these phagocytic cells in order to make them more efficient in fighting pathogenic organisms (Radostits et al., 2000).

It was also observed that dietary supplementation of HPLP resulted in a higher total protein (5.12 - 7.41 g/dL), albumin (3.09 - 4.42 g/dL) and 2.03 - 2.99 g/dL and values were within the bench mark reported by Mairbäurl (2003). Decrease in total protein concentration could be as a result of malnutrition, renal and liver dysfunction (Radostits et al., 2000). Albumin/globulin ratio range observed in this experiment (1.47 - 1.52 g/dL) was similar to the result of Alagbe (2024); Muhl and Liebert (2007a) who discovered that albumin/globulin ratio of piglets fed novel phytogenic feed additive varied from 1.20 - 1.72 g/dL. This result was higher than those presented by Muhl and Liebert (2007b) who found that albumin/globulin ratio of piglets fed phytogenics varied from 1.56 - 2.00 g/dL. Glucose values varied from 4.06 - 6.59 mmol/L was within the standard values recorded by Wilson et al. (1972). Glucose levels are mostly elevated during periods of malnutrition, stress due to environment/housing or poor management (Alagbe et al., 2013). Values of alanine aminotransferase, aspartate aminotransferase and alkaline phosphatase which varied from 76.87 - 79.88 (IU/L), 52.75 - 56.97 (IU/L) and 102.5 - 115.9 (IU/L) were all similar ($P>0.05$) in all the groups. Result obtained was in agreement with the baseline values [(alanine aminotransferase; 71.00 - 110.6 IU/L)], [(aspartate aminotransferase; 40.50 - 95.00 IU/L)] and [(alkaline phosphatase; 100.8 - 150.0 IU/L)] reported by Ventrella et al. (2017). This result suggests that dietary supplementation of HPLP at 120 g (G_E) did not compromise the integrity of the liver, kidney and other vital organs in the body of piglets.

Value of triglycerides, cholesterol, low density lipoprotein and high density lipoprotein took the form of 52.09 - 77.81 mg/dL, 100.1 to 131.6 mg/dL, 85.03 - 110.4 mg/dL and 45.88 to 67.05 mg/dL. However, values obtained within reference values; [(triglycerides; 47.11 - 80.90 mg/dL)], [(cholesterol; 100.1 to

131.6 mg/dL], [(high density lipoprotein; 45.88 to 67.05 mg/dL) and (low density lipoprotein; 85.03 to 110.4 mg/dL)] reported by Dubreuil and Lapierre (1997); Friedrichs et al. (2012). Cholesterol, triglycerides and low density lipoprotein were higher among piglets which received control diet while HPLC groups recorded low values. This result suggests that HPLP possess hypolipidemic properties, thus preventing coronary diseases in piglets (Singh et al., 2021). Conversely, high density lipoprotein was higher in G_D (90 g HPLP/kg diet) and G_E (120 g HPLP/kg diet), intermediate G_B (30 g HPLP/kg diet), G_C (60 g HPLP/kg diet) and lowest in G_A (control). High density lipoprotein is responsible for carrying cholesterol from the tissue to the liver where they are removed from the body of animals, thus preventing cardiovascular disease (Musa et al., 2020).

CONCLUSIONS

In conclusion, *Holarrhena pubescens* leaf powder contains numerous phyto-constituents which has medicinal properties. Supplementation of HPLP at 120 g per kg revealed the best result on growth performance. It was also able to suppress the activities of pathogenic organism and promote the proliferation of beneficial organisms in the gastrointestinal tract and strengthen the immune system without compromising the health status of piglets.

FURTHER STUDY

This research still has limitations, so further research is needed related to the topic Effect of dietary supplementation of *Holarrhena pubescens* leaf powder on the growth performance, carcass characteristics, microbial count and haemato-biochemical parameters of piglets in order to perfect this research and increase insight for readers.

REFERENCES

Jamadagni, P.S.; Pawar, S.D.; Jamadagni, S.B.; Chougule, S.; Gaidhani, S.N.; Murthy, S.N. (2017). Review of *Holarrhena antidysenterica* (L.) Wall. ex A. DC.: Pharmacognostic, Pharmacological, and Toxicological Perspective. *Pharmacogn. Rev.* 11, 141–144

Devi Prasad, A.G.; Shyma, T.B.; Raghavendra, M.P. (2013) Plants used by the tribes of for the treatment of digestive system disorders in Wayanad district, Kerala. *J. Appl. Pharm. Science*, 3: 171–175.

Dubey, D.; Padhy, R.N. (2012). Surveillance of multidrug resistance of two Gram-positive pathogenic bacteria in a teaching hospital and in vitro efficacy of 30 ethnomedicinal plants used by an aborigine of India. *Asian Pacific Journal of Tropical Disease*, 2: 273–281.

Ali, K.M.; Chatterjee, K.; De, D.; Jana, K.; Bera, T.K.; Ghosh, D. (2011). Inhibitory effect of hydro-methanolic extract of seed of *Holarrhena antidysenterica* on alpha-glucosidase activity and postprandial blood glucose level in normoglycemic rat. *Journal of Ethnopharmacology*, 135: 194–196.

Dua, V.K.; Gaurav, V.; Bikram, S.; Aswathy, R.; Upma, B.; Dayal, D.; Gupta, N.C.; Sandeep, K.; Ayushi, R. (2013). Anti-malarial property of steroid alkaloid conessine isolated from the bark of *Holarrhena antidysenterica*. *Malar. Journal*, 12: 194-196

Alagbe, J.O., Adeoye, Adekemi and Oluwatobi, O.A. (2020). Proximate and mineral analysis of *Delonix regia* leaves and roots. *International Journal on Integrated Education*. 3(10): 144-149.

Alagbe, J.O., Sharma, R., Eunice Abidemi Ojo, Shittu, M.D and Bello Kamoru Atanda (2020). Chemical evaluation of the proximate, minerals, vitamins and phytochemical analysis of *Daniellia oliveri* stem bark. *International Journal of Biological, Physical and Chemical Studies*, 2(1):16-22.

Maenner, K., Vahjen, W., Simon, O. (2011). Studies on the effects of essential-oil-based feed additives on performance, ileal nutrient digestibility, and selected bacterial groups in the gastrointestinal tract of piglets. *J. Anim. Sci.*, 89: 2106–2112.

Yan L and Kim I.H. (2012). Effect of eugenol and cinnamaldehyde on the growth performance, nutrient digestibility, blood characteristics, fecal microbial shedding and fecal noxious gas content in growing pigs. Asian-Aust. J. Anim. Sci., 25: 1178–1183

LiP., Piao, X., Ru, Y., Han, X., Xue, L., Zhang, H. (2012 a). Effects of adding essential oil to the diet of weaned pigs on performance, nutrient utilization, immune response and intestinal health. Asian-Australas. J. Anim. Sci., 25: 1617–1626.

Labaque, M.C., Kembro, J.M., Luna, A., Marin, R.H. (2013). Effects of thymol feed supplementation on female Japanese quail (*Coturnix coturnix*) behavioral fear response. Anim. Feed Sci. Technol., 183: 67–72.

Jacela, J.Y., DeRouchey, J.M., Tokach, M.D., Goodband, R.D., Nelssen, J.L., Renter, D.G., Dritz, S.S. (2010). Feed additives for swine: Fact sheets – prebiotics and probiotics, and phytogenics. J. Swine Health Prod., 18: 132–136

Kalimuthu, K., Arunprasath, A., 2018. Phytochemical and antioxidant activity of *Holarrhena pubescens*. International Journal of Pharmaceutical and Biological Achieves 9(2), 78-84.

Best, P., 2000. Health boosters from botany. Feed International. June, 15-16

Muhl, A and Liebert, F. (2007a). Growth and parameters of microflora in intestinal and fecal samples of piglets due to application of a phytogenic feed additive. J. Anim. Physiol. Anim. Nutr., 91: 411–418.

Muhl, A and Liebert, F. (2007b). No impact of a phytogenic feed additive on digestion and unspecific immune reaction in piglets. J. Anim. Physiol. Anim. Nutr., 91: 426–431.

Oetting, L. L., Utiyama, C.E., Giani, P.A., Ruiz, U.D., Miyada, V.S. (2006). Effects of herbal extracts and antimicrobials on apparent digestibility, performance, organs morphometry and intestinal histology of weanling pigs. Braz. J. Anim. Sci., 35: 1389–139

Pastorell, i.G., Rossi, R and Corino, C. (2012). Influence of *Lippia citriodora* verbascoside on growth performance, antioxidant status, and serum immunoglobulins content in piglets. *Czech J. Anim. Sci.*, 57: 312–322

Hanczakowska, E and Swiatkiewicz, M. (2012). Effect of herbal extracts on piglet performance and small intestinal epithelial villi. *Czech J. Anim. Sci.*, 57: 420–429.

Bartos, P., Dolan, A., Smutny L., Sístkova M., Celjal., Soch M., Havelka, Z. (2016). Effects of phytogenic feed additives on growth performance and on ammonia and greenhouse gases emissions in growing-finishing pigs. *Anim. Feed Sci. Technol.*, 212: 143–148

Bruno, D.G., Martins, S.M.M.K., Parazzi L. J., Afonso, E.R., DelSanto, T.A., Teixeira, S.D.N., Moreno, A.M., Moretti, A.D. (2013). Phytogenic feed additives in piglets challenged with *Salmonella typhimurium*. *Rev. Bras. Zootecn.*, 42: 137–143.

Cho, J.H., Zhang, S and Kim, I.H. (2012). Effects of anti-diarrhoeal herbs on growth performance, nutrient digestibility, and meat quality in pigs. *Asian-Australas. J. Anim. Sci.*, 25: 1595–1604.

Czech, A., Kowalczyk E and Grela, E.R. (2009). The effect of an herbal extract used in pig fattening on the animals' performance and blood components. *Ann. UMCS-EE*, 27: 25–33.

DeLange, C.F.M., Pluske, J.R., Gong, J., Nyachoti, C.M. (2010). Strategic use of feed ingredients and feed additives to stimulate gut health and development in young pigs. *Livest. Sci.*, 134: 124–134.

Dubreuil, P., Lapierre, H. (1997). Biochemistry Reference value for lactating dairy cows, nursery sows and growing pigs. *Canadian Journal of Veterinary Research*, 61: 235-239.

Fredrichs, K.R., Harr, K.E., Freeman, K.P. (2012). ASVCP reference interval guidelines: Determination of de novo reference intervals in veterinary spices and other related topics. *Veterinary Clinical Pathology*, 41:441-453.

Wilson, G.D., Harvey, D.G., Snook, C.R. (1972). A review of factors affecting blood chemistry in Pigs. *Brazilian Veterinary Journal*, 128: 596-610.

Ventrelia, D., Dondi, F., Barone, F. (2017). The biomedical piglet establishing reference intervals for haematology and clinical chemistry parameters of two age groups with or without supplementation, *BMC Veterinary Research*, 13(1): 23 - 27.

Kabir, A.K.L.; Begum, M.M.; Islam, T., 2018. Study of Bioactivities of *Holarrhena pubescens* growing in Bangladesh. *Dhaka University Journal of Pharmaceutical Science* 17, 131-137.

Wills, R. B. H., Bone, K., Morgan, M., 2000. Herbal products: active constituents, mode of action and quality control. *Nutrition Research Review* 13, 47-77

Savithramma, N., Linga Rao, M. and Suhrulatha, D. 2011. —Screening of Medicinal Plants for Secondary Metabolites.|| *Middle-East Journal of Science Research* 8:579-584.

Evans, W. C., Trease G. E. and Evans, D. 2002. *Trease and Evans Pharmacognosy*. 15th edition. Edinburgh, Saunders: Pp 249 and 454.

Edeoga, H. O., Okwu, D. E. and Mbaebie, B. O. 2005. Phytochemical constituents of some Nigerian medicinal plants. *African Journal of Biotechnology* 47:685- 688

Shin, Y., Tamai, Y. and Terazawa, M. 2001. Chemical Constituents of *Inonotus obliquus* IV. - Triterpene and Steroids from Cultured Mycelia. *Eurasian Journal for Forest Research* 2:27-30.

Alagbe, J.O., 2024. Effect of *Limonium stocksii* leaf powder on the growth performance and intestinal microbial population of broiler chicks. *World Journal of Agriculture and Forestry Sciences*, 2(2), 36-42.

Ojediran, T.K., Emiola, I.A., Durojaye, V., John, A., 2024a. Proximate, vitamin and GC-MS profiling of *Kigella africana* powder. *Cerrado: Agricultural and Biological Research*, 1(1): 13-20.

Ojediran, T.K., Emiola, I.A., Durojaye, V., Alagbe, J.O., 2024b. Analysis of *Kigellia africana* fruit's powder antioxidant and phytochemical properties. *Brazilian Journal of Science*, 3(7): 38-49.

Daniel N. Anorue, Friday Ubong and Alagbe Olujimi John (2023). Investigating the effects of pawpaw (*Carica papaya*) essential oil dietary supplementation on the growth performance and carcass characteristics of broilers. Research in: *Agricultural and Veterinary Sciences*, 7(3): 164 - 174.

Singh Sharma., Alagbe Olujimi John., Liu Xing., Sharma Ram and Kumar Amita (2022). Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectroometry. *International Journal of Agriculture and Animal Production*, 2(6): 18-27.

Adewale, A.O., Alagbe, J.O., Adeoye, Adekemi. O. (2021). Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phytonic Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices. *International Journal of Orange Technologies*, 3(3): 1-12.

Shittu, M.D., Alagbe, J.O., Adejumo, D.O., Ademola, S.G., Abiola, A.O., Samson, B.O and Ushie, F.T. (2021). Productive Performance, Caeca Microbial Population and Immune-Modulatory Activity of Broiler Chicks Fed Different Levels *Sida Acuta* Leaf Extract in Replacement of Antibiotics. *Bioinformatics and Proteomics Open Access Journal* 5(1): 000143.

Musa, B., Alagbe, J.O., Adegbite Motunrade Betty, Omokore, E.A. (2020). Growth performance, caeca microbial population and immune response of broiler chicks fed aqueous extract of *Balanites aegyptiaca* and *Alchornea cordifolia* stem bark mixture. *United Journal for Research and Technology*, 2(2):13-21.

Muritala, Daniel Shittu., Alagbe, J.O., Ojebiyi, O.O., Ojediran, T.K and Rafiu, T.A. (2022). Growth performance and haematological and serum biochemical parameters of broiler chickens given varied concentrations of *Polyalthia longifolia* leaf extract in place of conventional antibiotics. *Animal Science and Genetics* 18(2): 57-71.

Singh, A.S., Alagbe, J.O., Sharma, S., Oluwafemi, R.A and Agubosi, O.C.P. (2021). Effect of dietary supplementation of melon (*Citrullus linatus*) seed oil on the growth performance and antioxidant status of growing rabbits. *Journal of Multidimensional Research and Reviews*, 2(1): 78-95.

NRC. 2012. Nutrient requirements in swine. Washington (DC): Natl. Acad. Press.

Alagbe Olujimi John., Anuore, Daniel Nnadozie., Shittu Muritala Daniel and Ramalan, Sadiq Mohammad (2023). Growth performance and physiological response of weaned pigs fed diet supplemented with novel a phytogenics. *Brazilian Journal of Science*, 3(1): 43-57.

Wang, J., Zhu, F., Zhou, X. M., Niu, C. Y. and Lei, C. L. 2006. Repellent and fumigant activity of essential oils from *Artemisia vulgaris* to *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Journal of Stored Products Research* 42:339-347.

Alagbe, J.O. (2021). *Prosopis africana* stem bark as an alternative to antibiotic feed additives in broiler chick's diets: Performance and Carcass characteristics. *Journal of Multidimensional Research and Reviews*, 2(1): 64-77.

Ugbogu, O. C. and Akukwe, A. R. 2009. The antimicrobial effect of oils from *Pentaclethra macrophylla* Bent, *Chrysophyllum albidum* G.Don and *Persea gratissima* Gaerth F on some local clinical bacteria isolates. *African Journal of Biotechnology* 8.2:285-287

Alagbe, J.O., Adedeji, M.O., Habiba, Z., Nwosu, Gloria and Wyedia Dabara Comfort (2021). Physico-chemical properties of *Indigofera zollingeriana* seed oil. *Asian Journal of Advances in Medical Science* 3(4): 306-308.

John, A.O. (2024a). Growth performance, haemato-biochemical indices of broiler chicken fed *Aristolochia indica* as a phytogenic feed additive. *Cerrado: Agricultural and Biological Research*, 1(1), 42-53.

John, A.O. (2024b). *Clerodendron splendens* leaf extract supplementation in weaner rabbits: impact on growth performance, haematology and intestinal microbial population. *Cerrado: Agricultural and Biological Research*, 1(1), 21-31.

John, A.O. (2024c). Effect of coconut shell extract on the growth performance and some haemato-biochemical parameters of broiler chicken. *Brazilian Journal of Science*, 3(6): 82-95.

John, A.O. (2024d). Impact of dietary supplementation of *Rhamnus prinoides* leaf extract on the growth performance, nutrient retention and intestinal microbial count of Japanese quails. *Brazilian Journal of Science*, 3(5), 40-50.

John, A.O. (2024e). Effect of performance, serum biochemistry and

hematological components of feeding Japanese quails phytogenic feed additions comparising *Megaphrynum macrostachyum* leaves. *Brazilian Journal of Science*, 3(5), 51-64.

Tepe, B., Daferera, D., Sokmen, A., Sokmen, M. and Polissiou, M. 2005. Antimicrobial and antioxidant activities of the essential oil and various extracts of *Salvia tomentosa* Miller (Lamiaceae). *Food Chemistry* 90:333-340.

Taivini, T., Angelina, C. L., Christine, S. and Francis, C. 2001. Volatile compounds in fresh pulp of pineapple (*ananas comosus* [L.] Merr) from French Polynesia. *Journal of Essential Oil Research* 13:314-318.

Winnicka A., 2011: Reference values of basic laboratory tests in veterinary medicine. Wydawnictwo SGGW, Warszawa [in Polish].

Mayengbam P., Tolenkhomba T.C., 2015: Seasonal variation of hematobiochemical parameters in indigenous pig: Zovawk of Mizoram. *Vet. World* 8 (6): 732-737.

Alagbe, J.O. (2020). Performance, hematology and serum biochemical parameters of weaner rabbits fed different levels of fermented *Lagenaria breviflora* whole fruit extract. *Advances in Research and Reviews*, 2020, 1:5.

Egeli A.K., Framstad T., Morberg H., 1998: Clinical biochemistry, haematology and body weight in piglets. *Acta Vet. Scand.* 39: 381-393.

Faustini M., Munari E., Colombani C., Russo V., Maffeo G., Vigo D., 2000: Haematology and plasma biochemistry of stamboek prepubertal gilts in Italy. Reference values. *J. Vet. Med. A Physiol., Pathol., Clin. Med.* 47: 525-532.

Friendship R.M., Henry S.C., 1996: Cardiovascular system, haematology and clinical chemistry. In *Diseases of swine*. A.D. Leman, B.E. Straw, W.L. Mengeling, S. D'Allaire, D.J. Taylor (Eds). Iowa State Univ. Press, USA: 3-11.

Chmielowiec-Korzeniowska A., Babicz M., Pyrz M., 2008: Levels of hematological parameters of pigs over the fattening period. *Ann. Univ. Mariae Curie Skłodowska Sect. EE Zootech.* 3: 19-24.

Cooper C.A., Moraes L.E., Murray J.D., Owens S.D., 2014: Hematologic and biochemical reference intervals for specific pathogen free 6-week-old Hampshire-Yorkshire crossbred pigs. *J. Anim. Sci. Biotech.* 5: 1–5.

Czech A., Grela E.R., 2002: Effect of microbial phytase and formic acid supplementation sow diets on performance and hematological parameters of blood. *Ann. Anim. Sci., Suppl.*, 2: 201–205.

Czech A., Kowalcuk E., Grela E.R., 2009: The effect of a herbal extract used in pig fattening on the animals' performance and blood components. *Ann. Univ. Mariae Curie Skłodowska Sect. EE Zootech.* 2: 25–33.

Shittu, M.D., Alagbe, J.O., Alaba, O., Okanlawon, E.O., Adelakun, F.A., Emmanuel, E.O and Adejumo, D.O. (2024). Effect of ginger, garlic and Negro pepper on the gut microbes, gut histomorphometry and pathological assessment of selected organs of broiler chickens. *Association of Deans of Agriculture in Nigerian Universities*, 5: 105-121.

Alagbe Olujimi John (2024). Novel phytogenics' impact on weaned pigs growth performance, haematology and serum biochemical indicators. *Black Sea Journal of Agriculture*, 7(2): 82-89.

Mairbäurl, H. 2013. Red blood cells in sports: effects of exercise and training on oxygen supply by red blood cells. *Front. Physiol.* 4:332. doi:10.3389/fphys.2013.00332

Ramayo-Caldas, Y., E. Márml-Sánchez, M. Ballester, J. P. Sánchez, R. González-Prendes, M. Amills, and R. Quintanilla. 2019. Integrating genome-wide co-association and gene expression to identify putative regulators and predictors of feed efficiency in pigs. *Genet. Sel. Evol.* 51:48. doi: 10.1186/s12711-019-0490-

Chmielowiec-Korzeniowska A., Babicz M., Pyrz M., 2008: Levels of hematological parameters of pigs over the fattening period. *Ann. Univ. Mariae Curie Skłodowska Sect. EE Zootech.* 3: 19–24.

Egeli A.K., Framstad T., Morberg H., (1998). Clinical biochemistry, haematology and body weight in piglets. *Acta Vet. Scand.* 39: 381–393.